

Table 22. Percent Vegetative Cover by Species in Seasonal Wetland Habitat - Waterbody 17
1999 Annual Monitoring Report
Former Fort Ord, Monterey County, California

Code	Scientific Name	Common Name	Transect 1								Average Cover (%)	Transect 2									Average Cover (%)	Combined Average Cover (%)
			1 %	2 %	3 %	4 %	5 %	6 %	7 %	8 %		1 %	2 %	3 %	4 %	5 %	6 %	7 %	8 %	9 %		
AGEX	<i>Agrostis exarata</i>	western bent-grass			5		1	5		1.4											0.69	
AICA	<i>Aira caryophyllea</i> *	silvery hair-grass			3		2			0.6											0.31	
BAPI	<i>Baccharis pilularis</i>	coyote brush												20					5	2.8	1.39	
BRMI	<i>Briza minor</i> *	small quaking grass			3			5		1.0											0.50	
BRDI	<i>Bromus diandrus</i> *	ripgut grass							5	0.6											0.31	
BRHO	<i>Bromus hordeaceus</i> *	soft chess									5									0.6	0.28	
COBO	<i>Conyza bonariensis</i> *	South American conyza				2	5			0.9											0.44	
COCA	<i>Conyza canadensis</i>	horseweed	2							0.3											0.13	
CYER	<i>Cyperus erythrorhizos</i>	red-rooting cyperus	10	20		15			10	6.9											3.44	
DACA	<i>Danthonia californica</i>	California oat-grass								0.6											0.31	
ELMA	<i>Eleocharis macrostachya</i>	pale spike-rush										30		30						6.7	3.33	
EPCI	<i>Epilobium ciliatum</i>	California willow-herb			1					0.1					5				5	1.1	0.62	
GNLA	<i>Gnaphalium luteo-album</i> *	weedy cudweed	1							0.1											0.06	
GNST	<i>Gnaphalium stramineum</i>	cotton-batting plant		10	10					2.5											1.25	
JUBA	<i>Juncus balticus</i>	Baltic rush								2.5											1.25	
JUBU	<i>Juncus b. bufonius</i>	common toad rush	2		10		10	10	10	5.3											2.63	
JUEF	<i>Juncus offusus</i>	common rush										20		20						4.4	2.22	
JUPA	<i>Juncus patens</i>	spreading rush												95	75					18.9	9.44	
JUPH	<i>Juncus phaeocephalus</i>	brown-headed rush				25	20	10	65	15.0		20		10					10	4.4	9.72	
LOMU	<i>Lolium multiflorum</i> *	Italian ryegrass	3						5	1.0											0.50	
LYHY	<i>Lythrum hyssopifolium</i> *	grass poly			2					0.3											0.13	
PLCO	<i>Plantago coronopus</i> *	cut-leaved plantain	2				8			1.3	70									7.8	4.51	
POMA	<i>Polypogon maritimus</i> *	maritime beard-grass	13		2					1.9											0.94	
POMO	<i>Polypogon monspeliensis</i> *	rabbitfoot grass					15			1.9											0.94	
RUSA	<i>Rumex salicifolius</i>	willow dock			5					0.6				3						0.3	0.48	
SALA	<i>Salix laevigata</i>	red willow							50	18.8											9.38	
SCRO	<i>Scirpus robustus</i>	prarie bulrush											55		5					6.7	3.33	
SPMA	<i>Spergularia marina</i>	salt-marsh sand spurry					5			0.6											0.31	
TYLA	<i>Typha latifolia</i>	broad-leaved cat-tail		55	15	30				12.5			5				100	90	65	28.9	20.69	
VUBR	<i>Vulpia bromoides</i> *	six-week fescue							5	0.6											0.31	
VUMY	<i>Vulpia m. myuros</i> *	rat-tail fescue									5										0.28	
Total Vegetative Cover			33	85	53	80	61	30	140	135	77.1	80	70	63	60	100	100	100	90	85	83.1	80.12
BG	--	Bare Ground	70	15	47	20	40	70	10		34.0	20	15		60		10		20		13.9	23.94
DV	--	Dead Vegetation											15	40			10	20			9.4	4.72
OW	--	Opon Water																	10		1.1	0.56
Total Cover			103	100	100	100	101	100	150	135	111.1	100	100	103	120	100	120	120	100	105	107.6	109.3

*Non-native species