

**Table 24. 2000 Burn Area Intercept Distances and Percent Cover for Shrub Species,
Bare, and Vegetated Ground for Mature Habitat
2000 Annual Monitoring Report
Former Fort Ord, Monterey County, California**

| Code | Scientific Name | Transect Length (feet) Common name | Transect BA-1 166.7 | | Transect BA-2 159.1 | | Transect BA-4 168.9 | | Transect BA-6 164.4 | | Transect BA-7 163.5 | |
|----------------------------|------------------------------------|---------------------------------------|------------------------|--------------|------------------------|--------------|------------------------|--------------|------------------------|--------------|------------------------|--------------|
| | | | Intercept Distance | (%) Cover | Intercept Distance | (%) Cover | Intercept Distance | (%) Cover | Intercept Distance | (%) Cover | Intercept Distance | (%) Cover |
| ADFA | <i>Adenostoma fasciculatum</i> | chamise | 35.3 | 21.2 | 43.4 | 27.3 | 3.3 | 2.0 | 37.2 | 22.6 | 31.4 | 19.2 |
| ARPU | <i>Arctostaphylos pumila</i> | sandmat manzanita | 38.9 | 23.3 | | | 12.4 | 7.3 | 13.7 | 8.3 | 2.4 | 1.5 |
| ARTO | <i>Arctostaphylos t. tomentosa</i> | shaggy-barked manzanita | 72.8 | 43.7 | 94.9 | 59.6 | 135.4 | 80.2 | 87.3 | 53.1 | 138.2 | 84.5 |
| BAPI | <i>Baccharis pilularis</i> | coyote brush | | | 0.7 | 0.4 | | | | | | |
| CECR | <i>Ceanothus cuneatus rigidus</i> | Monterey ceanothus | 2.4 | 1.4 | 46.3 | 29.1 | 10.8 | 6.4 | 15.2 | 9.2 | 1.8 | 1.1 |
| CEDE | <i>Ceanothus dentatus</i> | dwarf ceanothus | | | | | | | | | | |
| ERER | <i>Ericameria ericoides</i> | mock heather | | | 5.2 | 3.3 | | | 0.4 | 0.2 | | |
| EREF | <i>Ericameria fasciculata</i> | Eastwood's goldenbush | | | | | | | | | | |
| ERCO | <i>Eriophyllum confertiflorum</i> | golden yarrow | | | | | | | | | | |
| Gael | <i>Garrya elliptica</i> | coast silk tassel | | | | | 0.4 | 0.2 | | | 2.9 | 1.8 |
| HEAR | <i>Heteromeles arbutifolia</i> | toyon | | | | | | | | | | |
| LOSC | <i>Lotus scoparius</i> | deerweed | 1.5 | 0.9 | | | | | | | | |
| LUCH | <i>Lupinus chamissionis</i> | silver beach lupine | | | | | | | | | | |
| MIAU | <i>Mimulus aurantiacus</i> | sticky monkey-flower | 4.1 | 2.5 | | | 20.8 | 12.3 | | | | |
| QUAG | <i>Quercus a. agrifolia</i> | coast live oak | | | | | | | | | | |
| RHCA | <i>Rhamnus c. californica</i> | California coffeeberry | | | | | | | | | | |
| SAME | <i>Salvia mellifera</i> | black sage | 13.3 | 8.0 | | | 21.5 | 12.7 | 11.7 | 7.1 | 4.0 | 2.4 |
| TODI | <i>Toxicodendron diversilobum</i> | poison oak | | | | | | | | | | |
| Total Species Cover | | | 168.3 | 101.0 | 190.5 | 119.7 | 204.6 | 121.1 | 165.5 | 100.7 | 180.7 | 110.5 |
| BG | -- | Bare Ground | 18.2 | 10.9 | 18.5 | 11.6 | 5.5 | 3.3 | 16.8 | 10.2 | 11.0 | 6.7 |
| DV | -- | Dead Vegetation | 6.1 | 3.7 | | | 4.1 | 2.4 | 10.2 | 6.2 | 6.9 | 4.2 |
| VG | -- | Vegetated Ground | | | | | 0.5 | 0.3 | | | 0.3 | 0.2 |
| Total Cover | | | 192.6 | 115.5 | 209.0 | 131.4 | 214.7 | 127.1 | 192.5 | 117.1 | 198.9 | 121.7 |

*Non-native species

**Table 24. 2000 Burn Area Intercept Distances and Percent Cover for Shrub Species,
Bare, and Vegetated Ground for Mature Habitat
2000 Annual Monitoring Report
Former Fort Ord, Monterey County, California**

| Code | Scientific Name | Transect Length (feet) Common name | Transect BA-9 155.8 | | Transect BA-20 158.1 | | Transect BD-1 166.7 | | Transect BD-2 145.2 | | Transect BD-4 142.3 | |
|----------------------------|------------------------------------|---------------------------------------|------------------------|--------------|-------------------------|--------------|------------------------|--------------|------------------------|--------------|------------------------|--------------|
| | | | Intercept Distance | (%) Cover | Intercept Distance | (%) Cover | Intercept Distance | (%) Cover | Intercept Distance | (%) Cover | Intercept Distance | (%) Cover |
| A DFA | <i>Adenostoma fasciculatum</i> | chamise | 8.4 | 5.4 | 37.5 | 23.7 | 51.9 | 31.1 | 9.0 | 6.2 | 11.1 | 7.8 |
| ARPU | <i>Arctostaphylos pumila</i> | sandmat manzanita | | | 28.2 | 17.8 | | | | | | |
| ARTO | <i>Arctostaphylos t. tomentosa</i> | shaggy-barked manzanita | 135.1 | 86.7 | 90.6 | 57.3 | 103.0 | 61.8 | 123.2 | 84.8 | 130.7 | 91.8 |
| BAPI | <i>Baccharis pilularis</i> | coyote brush | | | | | | | | | | |
| CECR | <i>Ceanothus cuneatus rigidus</i> | Monterey ceanothus | 0.6 | 0.4 | 1.1 | 0.7 | 2.3 | 1.4 | | | | |
| CEDE | <i>Ceanothus dentatus</i> | dwarf ceanothus | | | | | | | | | | |
| ERER | <i>Ericameria ericoides</i> | mock heather | | | | | | | | | | |
| EREF | <i>Ericameria fasciculata</i> | Eastwood's goldenbush | | | | | | | | | | |
| ERCO | <i>Eriophyllum confertiflorum</i> | golden yarrow | | | | | | | | | | |
| G AEL | <i>Garrya elliptica</i> | coast silk tassel | | | | | | | | | | |
| HEAR | <i>Heteromeles arbutifolia</i> | toyon | | | | | | | | | | |
| LOSC | <i>Lotus scoparius</i> | deerweed | | | | | | | | | | |
| LUCH | <i>Lupinus chamissionis</i> | silver beach lupine | | | | | | | | | | |
| MIAU | <i>Mimulus aurantiacus</i> | sticky monkey-flower | | | | | 5.5 | 3.3 | | | 0.6 | 0.4 |
| QUAG | <i>Quercus a. agrifolia</i> | coast live oak | | | | | | | | | | |
| RHCA | <i>Rhamnus c. californica</i> | California coffeeberry | | | | | | | | | | |
| SAME | <i>Salvia mellifera</i> | black sage | 28.8 | 18.5 | 7.5 | 4.7 | 0.7 | 0.4 | | | 8.6 | 6.0 |
| TODI | <i>Toxicodendron diversilobum</i> | poison oak | | | | | | | | | | |
| Total Species Cover | | | 172.9 | 111.0 | 164.9 | 104.3 | 163.4 | 98.0 | 132.2 | 91.0 | 151.0 | 106.1 |
| BG | -- | Bare Ground | 10.8 | 6.9 | 12.7 | 8.0 | 16.1 | 9.7 | 14.5 | 10.0 | 1.3 | 0.9 |
| DV | -- | Dead Vegetation | | | | | 7.2 | 4.3 | | | | |
| VG | -- | Vegetated Ground | | | | | | | | | | |
| Total Cover | | | 183.7 | 117.9 | 177.6 | 112.3 | 186.7 | 112.0 | 146.7 | 101.0 | 152.3 | 107.0 |

*Non-native species

**Table 24. 2000 Burn Area Intercept Distances and Percent Cover for Shrub Species,
Bare, and Vegetated Ground for Mature Habitat
2000 Annual Monitoring Report
Former Fort Ord, Monterey County, California**

| Code | Scientific Name | Transect Length (feet) Common name | Transect BD-5 158.3 | | Transect BD-6 156.4 | | Transect BD-7 154.2 | | Transect BD-8 162.5 | | Transect BE-5 154.3 | |
|----------------------------|------------------------------------|---------------------------------------|------------------------|--------------|------------------------|--------------|------------------------|--------------|------------------------|--------------|------------------------|--------------|
| | | | Intercept Distance | (%) Cover | Intercept Distance | (%) Cover | Intercept Distance | (%) Cover | Intercept Distance | (%) Cover | Intercept Distance | (%) Cover |
| ADFA | <i>Adenostoma fasciculatum</i> | chamise | 21.8 | 13.8 | 41.9 | 26.8 | 35.2 | 22.8 | 19.0 | 11.7 | 32.8 | 21.3 |
| ARPU | <i>Arctostaphylos pumila</i> | sandmat manzanita | 4.2 | 2.7 | 7.0 | 4.5 | | | 19.1 | 11.8 | 16.8 | 10.9 |
| ARTO | <i>Arctostaphylos t. tomentosa</i> | shaggy-barked manzanita | 127.3 | 80.4 | 66.3 | 42.4 | 117.7 | 76.3 | 71.1 | 43.8 | 91.6 | 59.4 |
| BAPI | <i>Baccharis pilularis</i> | coyote brush | | | 14.5 | 9.3 | 0.5 | 0.3 | 5.3 | 3.3 | | |
| CECR | <i>Ceanothus cuneatus rigidus</i> | Monterey ceanothus | 1.5 | 0.9 | 1.5 | 1.0 | 11.0 | 7.1 | 9.0 | 5.5 | 2.2 | 1.4 |
| CEDE | <i>Ceanothus dentatus</i> | dwarf ceanothus | | | 12.5 | 8.0 | 1.8 | 1.2 | 2.2 | | | |
| ERER | <i>Ericameria ericoides</i> | mock heather | | | | | | | 3.5 | 2.2 | | |
| EREF | <i>Ericameria fasciculata</i> | Eastwood's goldenbush | | | | | | | | | | |
| ERCO | <i>Eriophyllum confertiflorum</i> | golden yarrow | | | | | | | | | | |
| GAEL | <i>Garrya elliptica</i> | coast silk tassel | | | | | | | | | 8.9 | 5.8 |
| HEAR | <i>Heteromeles arbutifolia</i> | toyon | | | | | | | | | | |
| LOSC | <i>Lotus scoparius</i> | deerweed | | | 3.1 | 2.0 | | | | | | |
| LUCH | <i>Lupinus chamissionis</i> | silver beach lupine | | | | | | | | | | |
| MIAU | <i>Mimulus aurantiacus</i> | sticky monkey-flower | | | | | | | | | | |
| QUAG | <i>Quercus a. agrifolia</i> | coast live oak | | | | | | | | | | |
| RHCA | <i>Rhamnus c. californica</i> | California coffeeberry | | | | | | | 0.4 | 0.2 | | |
| SAME | <i>Salvia mellifera</i> | black sage | | | 11.4 | 7.3 | | | 1.0 | 0.6 | 3.1 | 2.0 |
| TODI | <i>Toxicodendron diversilobum</i> | poison oak | | | | | | | | | | |
| Total Species Cover | | | 154.8 | 97.8 | 158.2 | 101.2 | 166.2 | 107.8 | 130.6 | 79.0 | 155.4 | 100.7 |
| BG | -- | Bare Ground | 10.2 | 6.4 | 32.0 | 20.5 | 18.1 | 11.7 | 28.5 | 17.5 | 19.3 | 12.5 |
| DV | -- | Dead Vegetation | 1.9 | 1.2 | | | | | | | 1.8 | 1.2 |
| VG | -- | Vegetated Ground | | | | | | | 9.4 | 5.8 | | |
| Total Cover | | | 166.9 | 105.4 | 190.2 | 121.6 | 184.3 | 119.5 | 168.5 | 102.3 | 176.5 | 114.4 |

*Non-native species

**Table 24. 2000 Burn Area Intercept Distances and Percent Cover for Shrub Species,
Bare, and Vegetated Ground for Mature Habitat
2000 Annual Monitoring Report
Former Fort Ord, Monterey County, California**

| Code | Scientific Name | Transect Length (feet) Common name | Transect BE-22 158.1 | | Transect BE-24 149.2 | | Transect BF-1 177.4 | | Transect BF-2 160.5 | | Transect BF-3 143.2 | |
|----------------------------|------------------------------------|---------------------------------------|-------------------------|--------------|-------------------------|--------------|------------------------|--------------|------------------------|--------------|------------------------|--------------|
| | | | Intercept Distance | (%) Cover | Intercept Distance | (%) Cover | Intercept Distance | (%) Cover | Intercept Distance | (%) Cover | Intercept Distance | (%) Cover |
| ADFA | <i>Adenostoma fasciculatum</i> | chamise | 54.2 | 34.3 | 13.6 | 9.1 | 34.9 | 19.7 | 27.0 | 16.8 | 13.7 | 9.6 |
| ARPU | <i>Arctostaphylos pumila</i> | sandmat manzanita | 13.6 | 8.6 | 0.9 | 0.6 | | | | | 7.4 | 5.2 |
| ARTO | <i>Arctostaphylos t. tomentosa</i> | shaggy-barked manzanita | 61.2 | 38.7 | 107.4 | 72.0 | 124.4 | 70.1 | 112.5 | 70.1 | 119.6 | 83.5 |
| BAPI | <i>Baccharis pilularis</i> | coyote brush | | | | | 4.6 | 2.6 | 2.0 | 1.2 | 5.7 | 4.0 |
| CECR | <i>Ceanothus cuneatus rigidus</i> | Monterey ceanothus | 7.5 | 4.7 | 13.6 | 9.1 | | | | | 0.6 | 0.4 |
| CEDE | <i>Ceanothus dentatus</i> | dwarf ceanothus | 5.3 | 3.4 | 12.6 | 8.4 | | | | | | |
| ERER | <i>Ericameria ericoides</i> | mock heather | | | | | | | | | | |
| EREF | <i>Ericameria fasciculata</i> | Eastwood's goldenbush | | | | | | | | | | |
| ERCO | <i>Eriophyllum confertiflorum</i> | golden yarrow | | | | | | | | | | |
| GAEL | <i>Garrya elliptica</i> | coast silk tassel | | | 1.2 | 0.8 | | | | | | |
| HEAR | <i>Heteromeles arbutifolia</i> | toyon | | | | | | | | | | |
| LOSC | <i>Lotus scoparius</i> | deerweed | | | | | | | | | | |
| LUCH | <i>Lupinus chamissionis</i> | silver beach lupine | | | | | | | | | | |
| MIAU | <i>Mimulus aurantiacus</i> | sticky monkey-flower | 0.5 | 0.3 | | | | | 9.5 | 5.9 | 2.0 | 1.4 |
| QUAG | <i>Quercus a. agrifolia</i> | coast live oak | 1.8 | 1.1 | | | | | | | | |
| RHCA | <i>Rhamnus c. californica</i> | California coffeeberry | | | | | | | | | | |
| SAME | <i>Salvia mellifera</i> | black sage | 6.4 | 4.0 | | | 7.1 | 4.0 | 17.5 | 10.9 | 15.2 | 10.6 |
| TODI | <i>Toxicodendron diversilobum</i> | poison oak | 7.6 | 4.8 | | | | | 0.0 | | 0.0 | |
| Total Species Cover | | | 158.1 | 100.0 | 149.3 | 100.1 | 171.0 | 96.4 | 168.5 | 105.0 | 164.2 | 114.7 |
| BG | -- | Bare Ground | 27.9 | 17.6 | 13.4 | 9.0 | 2.0 | 1.1 | 13.7 | 8.5 | 5.7 | 4.0 |
| DV | -- | Dead Vegetation | 2.6 | 1.6 | | | 4.4 | 2.5 | | | | |
| VG | -- | Vegetated Ground | | | | | | | | | | |
| Total Cover | | | 188.6 | 119.3 | 162.7 | 109.0 | 177.4 | 100.0 | 182.2 | 113.5 | 169.9 | 118.6 |

*Non-native species

**Table 24. 2000 Burn Area Intercept Distances and Percent Cover for Shrub Species,
Bare, and Vegetated Ground for Mature Habitat
2000 Annual Monitoring Report
Former Fort Ord, Monterey County, California**

| Code | Scientific Name | Transect Length (feet) Common name | Transect BF-4 170.8 | | Transect BF-5 147.4 | | Transect BG-2 158.8 | | Transect BG-3 154.9 | | Transect BG-5 143.7 | |
|---------------------|------------------------------------|---------------------------------------|------------------------|--------------|------------------------|--------------|------------------------|--------------|------------------------|--------------|------------------------|--------------|
| | | | Intercept Distance | (%) Cover | Intercept Distance | (%) Cover | Intercept Distance | (%) Cover | Intercept Distance | (%) Cover | Intercept Distance | (%) Cover |
| ADFA | <i>Adenostoma fasciculatum</i> | chamise | 12.0 | 7.0 | 13.8 | 9.4 | 80.9 | 50.9 | 18.0 | 11.6 | 12.9 | 9.0 |
| ARPU | <i>Arctostaphylos pumila</i> | sandmat manzanita | | | 29.9 | 20.3 | 17.6 | 11.1 | | | | |
| ARTO | <i>Arctostaphylos t. tomentosa</i> | shaggy-barked manzanita | 121.6 | 71.2 | 94.3 | 64.0 | 39.7 | 25.0 | 94.9 | 61.3 | 113.6 | 79.1 |
| BAPI | <i>Baccharis pilularis</i> | coyote brush | | | 16.0 | 10.9 | 5.9 | 3.7 | 0.4 | 0.3 | 0.5 | 0.3 |
| CECR | <i>Ceanothus cuneatus rigidus</i> | Monterey ceanothus | | | 27.9 | 18.9 | 18.4 | 11.6 | 5.1 | 3.3 | | |
| CEDE | <i>Ceanothus dentatus</i> | dwarf ceanothus | | | | | | | 2.1 | 1.4 | | |
| ERER | <i>Ericameria ericoides</i> | mock heather | | | | | 0.4 | 0.3 | | | | |
| EREF | <i>Ericameria fasciculata</i> | Eastwood's goldenbush | | | | | | | | | | |
| ERCO | <i>Eriophyllum confertiflorum</i> | golden yarrow | | | | | | | | | | |
| GAEL | <i>Garrya elliptica</i> | coast silk tassel | | | | | | | | | | |
| HEAR | <i>Heteromeles arbutifolia</i> | toyon | | | | | | | | | | |
| LOSC | <i>Lotus scoparius</i> | deerweed | | | | | | | | | | |
| LUCH | <i>Lupinus chamissionis</i> | silver beach lupine | | | | | | | | | | |
| MIAU | <i>Mimulus aurantiacus</i> | sticky monkey-flower | 6.0 | 3.5 | | | | | | | 15.5 | 10.8 |
| QUAG | <i>Quercus a. agrifolia</i> | coast live oak | | | | | | | | | | |
| RHCA | <i>Rhamnus c. californica</i> | California coffeeberry | | | | | | | | | | |
| SAME | <i>Salvia mellifera</i> | black sage | 52.9 | 31.0 | | | | | 37.9 | 24.5 | 13.2 | 9.2 |
| TODI | <i>Toxicodendron diversilobum</i> | poison oak | | | | | | | | | | |
| Total Species Cover | | | 192.5 | 112.7 | 181.9 | 123.4 | 162.9 | 102.6 | 158.4 | 102.3 | 155.7 | 108.4 |
| BG | -- | Bare Ground | 1.7 | 1.0 | 9.0 | 6.1 | 18.2 | 11.5 | 10.6 | 6.8 | 8.3 | 5.8 |
| DV | -- | Dead Vegetation | | | 5.7 | 3.9 | 2.1 | 1.3 | 5.9 | 3.8 | 8.3 | 5.8 |
| VG | -- | Vegetated Ground | 1.6 | 0.9 | 0.7 | 0.5 | | | | | | |
| Total Cover | | | 195.8 | 114.6 | 197.3 | 133.9 | 183.2 | 115.4 | 174.9 | 112.9 | 172.3 | 119.9 |

*Non-native species

**Table 24. 2000 Burn Area Intercept Distances and Percent Cover for Shrub Species,
Bare, and Vegetated Ground for Mature Habitat
2000 Annual Monitoring Report
Former Fort Ord, Monterey County, California**

| Code | Scientific Name | Transect Length (feet) Common name | Transect BG-7 164.7 | | Transect BG-8 154.4 | | Transect P1T-1 162.0 | | Transect P1T-2 183.0 | | Transect P2T-1 164.8 | |
|----------------------------|------------------------------------|---------------------------------------|------------------------|--------------|------------------------|--------------|-------------------------|--------------|-------------------------|--------------|-------------------------|--------------|
| | | | Intercept Distance | (%) Cover | Intercept Distance | (%) Cover | Intercept Distance | (%) Cover | Intercept Distance | (%) Cover | Intercept Distance | (%) Cover |
| ADFA | <i>Adenostoma fasciculatum</i> | chamise | 17.9 | 10.9 | 33.7 | 21.8 | 44.8 | 27.7 | 41.3 | 22.6 | 49.4 | 30.0 |
| ARPU | <i>Arctostaphylos pumila</i> | sandmat manzanita | 2.7 | 1.6 | | | 11.3 | 7.0 | 26.1 | 14.3 | 43.1 | 26.2 |
| ARTO | <i>Arctostaphylos t. tomentosa</i> | shaggy-barked manzanita | 106.5 | 64.7 | 51.7 | 33.5 | 57.2 | 35.3 | 102.6 | 56.1 | 52.4 | 31.8 |
| BAPI | <i>Baccharis pilularis</i> | coyote brush | | | 13.0 | 8.4 | | | | | | |
| CECR | <i>Ceanothus cuneatus rigidus</i> | Monterey ceanothus | 5.3 | 3.2 | 3.0 | 1.9 | 3.6 | 2.2 | 6.9 | 3.8 | 12.8 | 7.8 |
| CEDE | <i>Ceanothus dentatus</i> | dwarf ceanothus | 15.4 | 9.4 | 0.8 | 0.5 | | | | | 3.9 | 2.4 |
| ERER | <i>Ericameria ericoides</i> | mock heather | | | | | | | | | 5.3 | 3.2 |
| EREF | <i>Ericameria fasciculata</i> | Eastwood's goldenbush | | | | | 0.4 | 0.2 | | | | |
| ERCO | <i>Eriophyllum confertiflorum</i> | golden yarrow | | | | | 0.3 | 0.2 | | | 0.3 | 0.2 |
| GAEL | <i>Garrya elliptica</i> | coast silk tassel | | | | | | | 4.0 | 2.2 | | |
| HEAR | <i>Heteromeles arbutifolia</i> | toyon | | | | | | | | | | |
| LOSC | <i>Lotus scoparius</i> | deerweed | | | | | 0.7 | 0.4 | | | | |
| LUCH | <i>Lupinus chamissionis</i> | silver beach lupine | | | 3.1 | 2.0 | | | | | | |
| MIAU | <i>Mimulus aurantiacus</i> | sticky monkey-flower | | | 14.2 | 9.2 | 5.9 | 3.6 | 3.7 | 2.0 | | |
| QUAG | <i>Quercus a. agrifolia</i> | coast live oak | | | | | | | | | | |
| RHCA | <i>Rhamnus c. californica</i> | California coffeeberry | | | | | | | | | 2.0 | 1.2 |
| SAME | <i>Salvia mellifera</i> | black sage | 40.4 | 24.5 | 25.0 | 16.2 | 15.0 | 9.3 | 9.3 | 5.1 | | |
| TODI | <i>Toxicodendron diversilobum</i> | poison oak | | | 5.3 | 3.4 | | | | | | |
| Total Species Cover | | | 188.2 | 114.3 | 149.8 | 97.0 | 139.2 | 85.9 | 193.9 | 106.0 | 169.2 | 102.7 |
| BG | -- | Bare Ground | 12.6 | 7.7 | 44.0 | 28.5 | 31.1 | 19.2 | 20.2 | 11.0 | 20.0 | |
| DV | -- | Dead Vegetation | 2.2 | 1.3 | | | 1.9 | 1.2 | | | | |
| VG | -- | Vegetated Ground | 6.0 | 3.6 | 3.0 | 1.9 | | | 0.1 | 0.1 | 0.5 | 0.3 |
| Total Cover | | | 209.0 | 126.9 | 196.8 | 127.5 | 172.2 | 106.3 | 214.2 | 117.0 | 189.7 | 103.0 |

*Non-native species

**Table 24. 2000 Burn Area Intercept Distances and Percent Cover for Shrub Species,
Bare, and Vegetated Ground for Mature Habitat
2000 Annual Monitoring Report
Former Fort Ord, Monterey County, California**

| Code | Scientific Name | Transect Length (feet) Common name | Transect 16-1 165.9 | | Transect 16-2 163.0 | | Transect 17-1 155.7 | | Transect 17-2 163.4 | | Transect 18-1 156.1 | | Combined Estimated (%) Cover |
|----------------------------|------------------------------------|---------------------------------------|------------------------|--------------|------------------------|--------------|------------------------|--------------|------------------------|--------------|------------------------|--------------|---------------------------------------|
| | | | Intercept Distance | (%) Cover | Intercept Distance | (%) Cover | Intercept Distance | (%) Cover | Intercept Distance | (%) Cover | Intercept Distance | (%) Cover | |
| ADFA | <i>Adenostoma fasciculatum</i> | chamise | | | 8.6 | 5.3 | 2.6 | 1.7 | 8.3 | 5.1 | 0.9 | 0.6 | 16.29 |
| ARPU | <i>Arctostaphylos pumila</i> | sandmat manzanita | 59.5 | 35.9 | 40.2 | 24.7 | | | | | 14.0 | 9.0 | 7.34 |
| ARTO | <i>Arctostaphylos t. tomentosa</i> | shaggy-barked manzanita | 108.4 | 65.3 | 85.9 | 52.7 | 152.0 | 97.6 | 125.5 | 76.8 | 117.8 | 75.5 | 63.42 |
| BAPI | <i>Baccharis pilularis</i> | coyote brush | | | | | | | | | 1.1 | 0.7 | 1.26 |
| CECR | <i>Ceanothus cuneatus rigidus</i> | Monterey ceanothus | | | | | | | 5.9 | 3.6 | 3.6 | 2.3 | 3.95 |
| CEDE | <i>Ceanothus dentatus</i> | dwarf ceanothus | | | | | | | | | 0.9 | 0.6 | 1.03 |
| ERER | <i>Ericameria ericoides</i> | mock heather | | | | | | | | | | | 0.27 |
| EREF | <i>Ericameria fasciculata</i> | Eastwood's goldenbush | | | | | | | | | | | 0.01 |
| ERCO | <i>Eriophyllum confertiflorum</i> | golden yarrow | | | 0.2 | 0.1 | | | 1.8 | 1.1 | 1.0 | 0.6 | 0.06 |
| GAEL | <i>Garrya elliptica</i> | coast silk tassel | | | 4.8 | 2.9 | | | | | | | 0.40 |
| HEAR | <i>Heteromeles arbutifolia</i> | toyon | | | 1.6 | | | | | | | | 0.03 |
| LOSC | <i>Lotus scoparius</i> | deerweed | | | | | | | | | | | 0.10 |
| LUCH | <i>Lupinus chamissionis</i> | silver beach lupine | | | | | | | | | | | 0.06 |
| MIAU | <i>Mimulus aurantiacus</i> | sticky monkey-flower | | | | | 0.3 | 0.2 | 3.7 | 2.3 | 0.8 | 0.5 | 1.67 |
| QUAG | <i>Quercus a. agrifolia</i> | coast live oak | | | | | | | 1.0 | 0.6 | | | 0.05 |
| RHCA | <i>Rhamnus c. californica</i> | California coffeeberry | | | | | | | | | | | 0.04 |
| SAME | <i>Salvia mellifera</i> | black sage | | | 18.3 | 11.2 | | | 33.6 | 20.6 | | | 7.24 |
| TODI | <i>Toxicodendron diversilobum</i> | poison oak | | | | | | | | | 15.1 | 9.7 | 0.50 |
| Total Species Cover | | | 167.9 | 101.2 | 159.6 | 96.9 | 154.9 | 99.5 | 179.8 | 110.0 | 155.2 | 99.4 | 103.70 |
| BG | -- | Bare Ground | 1.1 | 0.7 | 16.9 | 10.4 | 2.0 | 1.3 | 11.5 | 7.0 | | | 9.03 |
| DV | -- | Dead Vegetation | | | | | | | | | | | 1.28 |
| VG | -- | Vegetated Ground | | | | | | | | | 11.9 | 7.6 | 0.61 |
| Total Cover | | | 169.0 | 101.9 | 176.5 | 107.3 | 156.9 | 100.8 | 191.3 | 117.1 | 167.1 | 107.0 | 114.62 |

*Non-native species